

Prompt Optical Observations of Gamma-ray Bursts

Carl Akerlof¹, Richard Balsano², Scott Barthelmy³, Jeff Bloch², Paul Butterworth³, Don Casperson², Tom Cline³, Sandra Fletcher², Fillippo Frontera⁴, Galen Gisler², John Heise⁵, Jack Hills², Kevin Hurley⁶, Robert Kehoe¹, Brian Lee^{1,9}, Stuart Marshall⁷, Tim McKay¹, Andrew Pawl¹, Luigi Piro⁸, John Szymanski² and Jim Wren²

¹*University of Michigan, Ann Arbor, MI 48109*

²*Los Alamos National Laboratory, Los Alamos, NM 87545*

³*NASA/Goddard Space Flight Center, Greenbelt, MD 20771*

⁴*Università degli Studi di Ferrara, Ferrara, Italy*

⁵*Space Research Organization, Utrecht, The Netherlands*

⁶*Space Sciences Laboratory, University of California, Berkeley, CA 94720-7450*

⁷*Lawrence Livermore National Laboratory, Livermore, CA 94550*

⁸*Istituto Astrofisica Spaziale, Rome, Italy*

⁹*Fermi National Accelerator Laboratory, Batavia, IL 60510*

ABSTRACT

The Robotic Optical Transient Search Experiment (ROTSE) seeks to measure simultaneous and early afterglow optical emission from gamma-ray bursts (GRBs). A search for optical counterparts to six GRBs with localization errors of 1 square degree or better produced no detections. The earliest limiting sensitivity is $m_{ROTSE} > 13.1$ at 10.85 seconds (5 second exposure) after the gamma-ray rise, and the best limit is $m_{ROTSE} > 16.0$ at 62 minutes (897 second exposure). These are the most stringent limits obtained for GRB optical counterpart brightness in the first hour after the burst. Consideration of the gamma-ray fluence and peak flux for these bursts and for GRB990123 indicates that there is not a strong positive correlation between optical flux and gamma-ray emission.

Subject headings: gamma rays: bursts, observations

1. Introduction

The gamma-ray emission of GRBs typically has a duration of the order of tens of seconds or less and exhibits little pattern in its very pronounced temporal variation. Until 1997, the brevity of bursts prevented multi-wavelength observation which would allow an accurate localization of the source. As a result, the burst mechanism, environment, location and energy scale have been elusive. Since 1997, multi-wavelength afterglow observations (eg. Costa et al. (1997), van Paradjis et al (1997)) have established the distance to several bursts (eg. Metzger et al. (1997), Kulkarni et al. (1998)) and illuminated the physics processes occurring a few hours to days after the gamma-ray

onset. The burst mechanism, however, remains a mystery.

Prompt radiation provides critical detail about the processes of the burst itself. Detection of such emission, in the optical for instance, requires instruments with wide field-of-view, rapid response and automated operation. The ROTSE-I CCD telephoto array meets these objectives (see Kehoe et al. (1999)). Preliminary data from the BATSE detectors (Fishman et al. (1989)) on-board the Compton Gamma-Ray Observatory are used by the GRB Coordinates Network (GCN, Barthelmy et al. (1998), Barthelmy et al. (1995)) to generate triggers about once per day providing rough coordinates ($\Delta\theta \sim 10^\circ$) within ~ 5 seconds of a burst. A dynamic response points ROTSE-I at

the trigger coordinates within 3 seconds of their receipt. The discovery of prompt optical emission from GRB990123 (Akerlof et al. (1999)) has illustrated properties of early shock development and the immediate environment of bursts. This paper presents a further search for optical counterparts in a subset of our GRB trigger data, as well as a comparison of the results with GRB990123.

2. Observations and Reduction

The subset of 6 triggers discussed here were taken in the first year of operation. They were selected for this analysis because they possess localization errors of about 1 square degree or smaller (see Table 1). This positional accuracy, which reduces the search area and background by a factor of more than 200 from that available from BATSE alone, is generally obtained from the relative timing of signals from BATSE and the gamma-ray detectors on-board Ulysses (Hurley et al. (1999b)). Thin 'Interplanetary Network' (IPN) annuli are generated which are only about 0.1° wide (Hurley et al. (1999a)). The intersection of the BATSE position probability distribution with such a timing annulus produces an IPN arc a few degrees long. If available, a third detection reduces this arc to a smaller diamond-shaped region. In the current sample, 4 bursts are localized to IPN arcs, and GRB981121 has an IPN diamond using NEAR data. The BeppoSAX (Feroci et al. (1997), Jager et al. (1997)) satellite observed GRB980329 (Frontera et al. (1998)), providing a very accurate position. This sample does not include GRBs on the faint end of the BATSE fluence distribution or short bursts (see Kouveliotou et al. (1993), Kouveliotou et al. (1996)). Both limitations will be addressed in later analyses.

For prompt GRB triggers, we initially begin taking 5 second exposures to retain sensitivity to rapid variation, then lengthen to 25 second and 125 second exposures to maximize sensitivity. If the trigger position error is of the same order as the ROTSE-I field-of-view ($16^\circ \times 16^\circ$), we also 'tile' around the given position at specific epochs in the sequence to ensure coverage of sources with errant initial positions but well-localized later. We then return to the direct pointing with longer exposures and begin the sequence again.

ROTSE-I first triggered on GRB980329 and be-

gan the first exposures 11.5 seconds after the burst had started. Unfortunately, the sky was cloudy for the early data and hazy for the later images. Nevertheless, some early images are clear enough in the immediate region of the burst to detect 10th magnitude objects. To maximize sensitivity in later, clearer images, the last two observations are the result of co-adding two and three frames, respectively. GRB980401 occurred during focusing tests so a manual response of eight exposures was performed. The last four 125 second exposures were co-added into one observation spread over 897 seconds. The final localization for GRB980420 places it near the galactic plane (most probable $g_b \sim -10^\circ$), and focused images are very crowded. The optics of the two cameras covering the main part of the IPN arc, however, were poorly focused. The final localization for GRB980627 places a majority of the probable area outside of even tiled exposures with the result that we have 40% coverage in only four tiled images. Observing conditions for GRB981121 and GRB981223 were good.

Raw images are dark subtracted and flat-fielded, followed by source finding in the corrected images using SExtractor (Bertin and Arnouts (1996)). We then perform an astrometric and photometric calibration by comparison to the Hipparcos catalog (Høg et al. (1998)). Our astrometric errors are 1.4 arcsec. Since we operate with unfiltered CCDs to maximize light-gathering ability, photometry is established by comparing raw ROTSE magnitudes to V-band measures and color correcting based on $B - V$. The resulting magnitude, m_{ROTSE} , corresponds on average to m_V but includes sensitivity in the B, V, I and especially R bands. Our photometric errors are 0.02 magnitude for stars brighter than magnitude 12.

3. Analysis and Discussion

Due to observation of an X-ray counterpart for GRB980329, optical (Djorgovski et al. (1998), Palazzi et al. (1998), Pedersen et al. (1998b)) and radio (Taylor et al. (1998)) counterparts were observed several hours later. For such precise localizations, we would accept any detection at the known location of the burst. No optical emission was observed, so for the early images we take the sensitivity to be 0.5 magnitudes brighter than the

dimmiest SAO and GSC stars visible in the immediate region of the burst. We calculated the limiting sensitivities for the co-added frames by extrapolation of the Hipparcos-derived calibration to our 5σ threshold. A cross-check of this calibration was performed by directly comparing to the USNO catalog and finding the faintest m_{ROTSE} to which we are more than 50% efficient. Given the afterglow measurements, we are able to place a constraint on the overall power-law decline of optical emission from GRB980329 to be shallower than $t^{-1.8}$ with respect to the earliest afterglow detection. This contrasts with the faster decline of the X-ray emission (Greiner et al. (1998)).

For the other bursts, we require that a source magnitude vary by at least $0.5 + 5\sigma$ where σ is the statistical error on the dimmiest measurement. Varying objects are only considered bona fide optical counterpart candidates if they appear in at least two successive images. This removes backgrounds such as cosmic rays and satellite glints which show up frequently in ROTSE-I images. No rapidly varying objects were found in the allowed error regions of these five bursts. The image sensitivities were determined from the Hipparcos calibration as described above, and in most cases were cross-checked with the USNO catalog comparison.

Limiting magnitudes are given in Table 2 for up to three epochs marking significant improvements in sensitivity. Figure 1 displays all relevant observations where coverage exceeded 50%, and indicates that ROTSE-I has sensitivity to optical bursts significantly fainter than GRB990123. The earliest limit is $m_{ROTSE} > 13.1$ at 10.85 seconds for GRB981223. The best limit of this sample is $m_{ROTSE} > 16.0$ at 62 minutes for GRB980401. We can conclude that bright optical counterparts (ie. $m_{ROTSE} \sim 10$) are uncommon.

Since prompt optical emission has been seen in GRB990123, we ask whether optical emission from a GRB is correlated with gamma-ray output, as is suggested by Sari and Piran (1999). Because we do not know whether fluence or peak flux are accurate measures of the total gamma ray emission, we consider both in our comparison. To bring all bursts onto a common footing, we first adjust their m_{ROTSE} limits by $2.5 \log(f/f_{GRB990123})$, where f is the gamma-ray fluence. We calculate this fluence to be that measured in the BATSE 50 - 100 keV plus 100 - 300 keV channels to avoid system-

atics due to problems in spectral fitting the other channels (Briggs (1999)). These fluence-scaled limits are plotted along with the GRB990123 observations in Figure 2. We have also adjusted our optical limits by scaling according to the BATSE measure of peak flux in the 64ms binning of the 50 - 300 keV data (see Figure 3).

Variation of galactic extinction over the IPN arcs prevents us from quoting an accurate value for most of these bursts. However, it is much less than 1 magnitude at their most probable locations. Since GRB990123 has a similar low value of extinction ($= 0.04$), the effect of galactic extinction on our comparison should be minimal. The one exception is GRB980420 which may have over 2 magnitudes of extinction. Although extinction near the source can only be measured for GRB980329, it is likely most GRBs are not so heavily obscured since the great majority observed to have both X-ray and radio counterparts also exhibit optical emission (Frail (1999)).

Under the assumption of gamma-ray scaling, ROTSE-I is sensitive to GRB990123-like optical bursts for GRB981223 from 30 to 300 seconds. Around 1 minute, the optical emission of GRB981121 and GRB981223 would have been more than 2 magnitudes over our detection threshold. Either GRB990123 is atypical of GRBs in general, or there is not a strong correlation of optical flux with gamma-ray emission and the inherent dispersion to any actual correlation must be larger than two magnitudes to explain the results from GRB981121 and GRB981223.

4. Conclusions

In a study of six well-localized gamma-ray bursts, no optical counterparts were identified. When comparing to afterglow observations of GRB980329, we constrain the overall power-law decay of the optical emission to be shallower than $t^{-1.8}$. When using either gamma-ray fluence or peak flux as a predictor of optical emission, we find that especially around 1 minute optical emission is at least two magnitudes dimmer than for GRB990123. This non-detection of another optical burst indicates that optical emission is not strongly correlated with gamma-ray output.

We thank the BATSE team for their GRB data and Michael Briggs for further assistance. We also thank the NEAR team for their GRB981121 data. ROTSE is supported by NASA under *SR&T* grant NAG5-5101, the NSF under grants AST-9703282 and AST-9970818, the Research Corporation, the University of Michigan, and the Planetary Society. Work performed at LANL is supported by the DOE under contract W-7405-ENG-36. Work performed at LLNL is supported by the DOE under contract W-7405-ENG-48.

REFERENCES

- AKERLOF, C. ET AL. 1999. *Nature*. **398**:400–402.
- BARTHELMY, S. ET AL. 1995. *Astrophysics and Space Science*, **231**:235–238.
- BARTHELMY, S. ET AL. 1998. <http://gcn.gsfc.nasa.gov/gcn/>.
- BERTIN, E. AND ARNOUTS, S. 1996. *Astron. Astrophys. Suppl.*, **117**:393–404.
- BRIGGS, M. 1999. private communication.
- COSTA, E. ET AL. 1997. *Nature*. **387**:783–785.
- DJORGovski, S. ET AL. 1998. *GCN Circ.* **41**.
- FEROCI, M. ET AL. 1997. In Proc. SPIE Vol. 3114, *EUV, X-ray, and Gamma-Ray Instrumentation for Astronomy VIII*, 186+.
- FISHMAN, G. ET AL. 1989. In W. Neil Johnson, ed., *Proc. of the Gamma Ray Observatory Science Workshop*, 2–39.
- FRAIL, D. ET AL. 1999. To appear in R. Kippen, R. Mellozzi, V. Connaughton, ed., *Gamma-Ray Bursts 5th Huntsville Symposium*, astro-ph/9912171.
- FRONTERA, F. ET AL. 1998. *IAU Circ.* **6853**.
- GREINER, J. ET AL. 1998. *GCN Circ.* **59**.
- HØG, E. ET AL. 1998. *Astron. Astrophys.* **335**:L65–L68.
- HURLEY, K. ET AL. 1999. *ApJ Suppl.* **120**:399–408.
- HURLEY, K. ET AL. 1999. *ApJ Suppl.* **122**:497.
- JAGER, R. ET AL. 1997. *Astron. Astrophys. Suppl.* **125**:557–572.
- KEHOE, R. ET AL. 1999. *STScI May Symp.*
- KOUVELIOTOU, C. ET AL. 1993. *ApJ* **413**:L101–L104.
- KOUVELIOTOU, C. ET AL. 1996. In C. Kouveliotou, M. Briggs, and G. Fishman, ed., *Gamma-Ray Bursts 3rd Huntsville Symposium*, 42–46.
- KULKARNI, S. ET AL. 1998. *Nature*, **393**:35–39.
- METZGER ET AL. 1997. *IAU Circ.* **6676**.
- PALAZZI, E. ET AL. 1998. *GCN Circ.* **48**.
- PEDERSEN, H. ET AL. 1998. *GCN Circ.* **52**.
- SARI, R. AND PIRAN, T. 1999. *ApJ*. **517**(2):L109–L112.
- TAYLOR, G. ET AL. 1998. *GCN Circ.* **40**.
- VAN PARADJIS, J. ET AL. 1997. *Nature*. **386**:686–689.

Fig. 1.— m_{ROTSE} limiting magnitudes vs. time after gamma-ray onset. GRB990123 optical burst detections are shown for comparison.

Fig. 2.— Limits rescaled by fluence for five GRBs vs. time after gamma-ray onset. If optical emission were positively correlated with gamma-ray fluence, ROTSE-I would have detected optical bursts for GRB981121 and GRB981223.

Fig. 3.— Flux rescaled limits for five GRBs vs. time after gamma-ray onset. If optical emission were positively correlated with peak gamma-ray flux, ROTSE-I would have detected optical bursts for GRB981121 and GRB981223.

date	trigger	T_{90}	fluence	ϕ_{peak}	source	coverage	t_+
980329	6665	19	322	13.8	SAX	100 (100)	11.5
980401	6672	28	40	5.60	IPN	87	2478
980420	6694	40	77	4.28	IPN	90 (96)	28.99
980627	6880	11	7	1.27	IPN	- (40)	11.97
981121	7219	54	70	1.33	IPN	100 (100)	14.51
981223	7277	30	128	7.44	IPN	100 (67)	8.35
990123	7343	63	1020	17.0	SAX	100 (100)	22.18

Table 1: Characteristics of six bursts responded to by ROTSE-I. For comparison, corresponding information for GRB990123 is also given (Akerlof et al. (1999)). The columns specify: GRB date, BATSE trigger number, duration in seconds, fluence ($\times 10^{-7} \text{erg/cm}^2$), peak flux (ϕ_{peak} , in $\text{photons/cm}^2/\text{s}$), source of best gamma-ray localization (see text), coverage of the GRB probability (%), and start time (t_+ , in sec.) for first image recorded. Coverages for tiled epochs are indicated in parentheses.

date	t_1	Δt_1	$m_{ROTSE}(t_1)$	t_2	Δt_2	$m_{ROTSE}(t_2)$	t_3	Δt_3	$m_{ROTSE}(t_3)$
980329	51.5	5	9.5	416	25	10.5	2239	390	14.7
980401	-	-	-	2485	15	14.1	3726	897	16.0
980420	31.49	5	11.1	176.32	5	12.4	578.21	25	12.5
980627	-	-	-	180.49	5	13.2	601.87	25	13.6
981121	17.01	5	13.3	219.28	25	14.9	742.05	125	15.3
981223	10.85	5	13.1	90.13	5	14.7	736.43	75	15.3

Table 2: Summary of limits for six bursts responded to by ROTSE-I. Columns list up to three epochs (middle of exposure, in sec.), and their exposure length (sec.) and sensitivity.





